

Variability in response to albuminuria lowering drugs: true or random?

S.I. Petrykiv¹, D de Zeeuw¹, F Persson², P Rossing^{2,3,4}, R.T. Gansevoort⁵, G.D. Laverman⁶,
H.J.L. Heerspink¹

¹Department of Clinical Pharmacy and Pharmacology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

²Steno Diabetes Center, Gentofte, Denmark

³NNF Center for Basic and Metabolic Research, Copenhagen University, Denmark

⁴HEALTH, Aarhus University, Aarhus, Denmark

⁵Department of Nephrology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

⁶Ziekenhuisgroep Twente, Almelo, the Netherlands

Correspondence to:

Hiddo J.L. Heerspink PhD

Department of Clinical Pharmacy and Pharmacology

University Medical Center Groningen

Postbox 30001

9700 RB Groningen

The Netherlands

Tel: +31 50 3617589

Fax: +31 50 3617889

e-mail: h.j.lambers.heerspink@umcg.nl

Running head: Variability in drug response: true or random?

Keywords: Drug response, albuminuria, precision medicine, personalized medicine

Word count: 2359

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/bcp.13217

Word count abstract: 250

Example of Tables of Links

Please insert these tables and the paragraph shown below the tables into your main manuscript document,

LIGANDS
Aliskiren
Atrasentan
Enalapril
Fosinopril
Irbesartan
Losartan
Paricalcitol
TARGETS
AT2 receptor
Endothelin receptor
Vitamine D receptor
Enzymese
Angiotensin converting enzyme
Renin

These Tables of Links list key protein targets and ligands in this article that are hyperlinked to corresponding entries in <http://www.guidetopharmacology.org>, the common portal for data from the IUPHAR/BPS Guide to PHARMACOLOGY (Southan et al., 2016), and are permanently archived in The Concise Guide to PHARMACOLOGY 2015/16 (^{a,b,c,d,e}Alexander et al., 2015a,b,c,d,e).*

Abstract

Aims: Albuminuria lowering drugs have shown different effect size in different individuals. Since urine albumin levels are known to vary considerably from day-to-day, we questioned whether the between-individual variability in albuminuria response after therapy initiation reflects a random variability or a true response variation to treatment. In addition, we questioned whether the response variability is drug dependent.

Methods: To determine whether the response to treatment is random or a true drug response, we correlated in six clinical trials the change in albuminuria during placebo or active treatment (on-treatment) with the change in albuminuria during wash-out (off-treatment). If these responses correlate during active treatment it suggests that at least part of the response variability can be attributed to drug response variability. We tested this for the following drugs: enalapril, losartan, aliskiren, atrasentan, and paricalcitol.

Results: No correlation between the on- and off-treatment albuminuria change was observed in the placebo arm of all clinical trials ($R^2 < 0.01$). However, we observed significant associations between the on- and off-treatment response (R^2 0.14 to 0.57; all $P < 0.015$) for different albuminuria lowering drugs. Additionally, the albuminuria responses strongly correlated when the same individual was re-exposed to the same drug at the same dose: lisinopril 10 mg/d ($R^2 = 53\%$; $P < 0.01$), losartan 50 mg/d ($R^2 = 63\%$; $P < 0.01$).

Conclusion: The degree of albuminuria lowering with anti-albuminuric drugs varies between patients. This variability in response appears drug-class independent. Identifying which factors determine this initial short-term variation in drug response appears important since the degree of albuminuria lowering is related to subsequent long-term renoprotection.

Introduction

Intervention in the Renin-Angiotensin-System (RAS) is the cornerstone of treatment to delay progression of renal disease in both diabetic and non-diabetic chronic kidney disease. The effects of RAS intervention on slowing renal function decline vary to a considerable extent between patients, ranging from a complete arrest of renal function loss to no effect.[1] This between-patient response variability on renal outcome is also observed in the short-term response in the renal risk markers including albuminuria and blood pressure. In fact, the initial change in albuminuria is in many cases a good indicator for the ultimate renal protective effect of the drug.[2, 3]

Changes in albuminuria over time, however, can also be the result of other factors not related to drug intervention. It is well known that albuminuria shows a substantial within-individual day-to-day variability which can be attributed to physical activity,[4, 5] analytical factors, measurement variability,[6] posture,[7, 8] or fever.[9] These daily fluctuations hamper the accuracy of drug efficacy monitoring. It has been suggested that short-term changes in risk markers can be largely attributed to background variability in the risk marker rather than individual drug response variability thereby limiting the value of these measures in individual drug response monitoring.[10] However, this suggestion is based on secondary analyses of large multi-centre trials and the conclusions may be biased by many factors such as changes in background medication over time, suboptimal drug adherence, and laboratory drift and variability over time.

To obtain more insight as to whether changes in albuminuria can be used to assess individual drug responsiveness unrelated to other factors, we analysed a series of clinical trials that tested the effects of different interventions on albuminuria. We first assessed the within-individual day-to-day variability in albuminuria in placebo arms of randomized controlled trials. Secondly, we determined whether the change in albuminuria following

treatment initiation correlates with the change in albuminuria following treatment discontinuation in active and placebo groups of randomized controlled trials. We postulate that if these changes in albuminuria correlate in the active treatment group but not in the placebo group, variable drug response is real. Thirdly, we measured albuminuria changes in patients who were re-challenged to the same drug at the same dose, to validate a true drug response variation. Finally, we tested whether these findings are consistent when different albuminuria lowering drugs were used.

Methods

Data sources

Day-to-day variability in albuminuria

To assess the day-to-day variability in albuminuria we calculated the intra-individual coefficient of variation in the placebo arms of PREVEND-IT, BENEDICT, IRMA-2, and RENAAL clinical trials.[11-14] The characteristics of the included populations are described elsewhere. In brief, the trials included patients with type 2 diabetes and non-diabetes with either normo-, micro-, or macroalbuminuria. First morning voided urines were collected in all trials, except PREVEND-IT in which 24-hour urine samples were collected, for assessment of albuminuria at baseline and every six months thereafter except in PREVEND in which albuminuria was measured at yearly intervals.

Between-individual drug response variability in albuminuria

To assess drug response variability in albuminuria we assessed the association in change of albuminuria directly after the initiation and cessation of drug treatment. To this end, we used the individual patient data of clinical trials that measured albuminuria in the first weeks after study treatment and drug discontinuation in six clinical studies.[11, 13, 15-18]. A systematic review of the literature was also conducted to identify additional studies. The search strategy and study flow diagram are described in supplement 1. Although a couple of additional studies were identified, we were unable to obtain the individual patient data. The study protocols of all included studies were approved by independent ethics committees. All participating patients signed informed consent before any study specific procedure commenced. Four studies tested the effect of RAS intervention, one the effect of an endothelin receptor antagonist, and another a vitamin D receptor antagonist. Four studies had a parallel design and two studies a cross-over design. Four studies had a follow-up ranging between four and 24 weeks whereas follow-up was two and four years in two other studies. Albuminuria was the primary surrogate end point in all trials. Albuminuria was measured 4 weeks after treatment discontinuation in all studies except in one study in which it was measured 12 weeks after treatment discontinuation. Albuminuria or proteinuria was measured in 24-hour urine collections in three studies, and in the remaining four studies the albumin to creatinine ratio was measured in first morning void urine collections.

We identified one clinical study to investigate the consistency in drug response exposing the same patient twice to the same dose of the same intervention. In that cross-over study non-diabetic proteinuric patients were exposed twice to six weeks treatment to either losartan 50 mg/day or lisinopril 10 mg/day .[19]

Statistical analysis

Continuous variables are described as mean and standard deviation, or median and 25th to 75th percentile. Categorical variables are presented as proportions or ratios. Albuminuria was log-transformed before analysis to take into account its non-parametric distribution and reported as geometric mean change. The within-individual coefficient of variation over time was calculated per individual as the standard deviation of three measurements divided by their geometric mean value. Kruskal Wallis was used with Dunn's post hoc test to determine statistical significance in the coefficient of variation in patients with normo-, micro-, or macroalbuminuria. Deming regression was applied to assess the association between individual changes in albuminuria after treatment initiation and discontinuation. To determine what proportion of the variability in albuminuria could be attributed to response variability, Deming regression was used to assess the correlation in albuminuria responses when the same individual was exposed twice to the same drug at the same dose. In addition, we calculated the positive predictive value, negative predictive value and area under the receiver operator characteristic for a 40% and 50% reduction in albuminuria during the second treatment period using the responses in albuminuria during the first treatment period with the same drug-dose combination as predictor. The 40% and 50% threshold were chosen since albuminuria decreased by 40% to 50% in this study. All analyses were conducted using StataSE version 13 for Windows. In all analyses, a P-value of less than 0.05 was used to reject the null hypothesis.

Results

Within individual variability in albuminuria over time

To assess the day-to-day within-individual variability in albuminuria (without any drug intervention), we firstly calculated the within-individual coefficient of variation (CV) in albuminuria using all samples over the full length of the placebo arm in four different trials.

In the overall population the median within-individual CV in albuminuria was 31.8% over a 12-month time period (range between trials 20.0 to 37.7%, Table 1). The within-individual CV was consistent regardless of the period during which it was assessed. The within-individual CV was somewhat higher in patients with micro- or macroalbuminuria at baseline compared to those with normoalbuminuria (Table 1 and Figure 1).

Correlation between the initial and post-treatment albuminuria change

Secondly, to assess whether the change in albuminuria after drug initiation correlates with the change after drug discontinuation, we analyzed six randomized placebo-controlled trials that assessed the effects of various drugs on albuminuria. The baseline characteristics of the trials are reported in Table 2. In all trials active treatment significantly decreased albuminuria with mean reductions ranging from 19% to 59% (Table 3). Across all trials, the mean decrease in albuminuria after treatment initiation significantly correlated with the mean increase following treatment discontinuation ($R^2=0.907$; $P<0.01$; Figure 2). Additionally, within each trial, the change in albuminuria following treatment initiation significantly correlated with the change in albuminuria after treatment discontinuation in the active treatment arms of each study except in the case of paricalcitol treatment (Table 3). These correlations remained statistically significant after adjustment for baseline blood pressure or changes in blood pressure. There was no statistically significant correlation in the placebo arms of all trials. In two additional trials in which patients with high normo- or microalbuminuria were followed

for 2 and 4 years before treatment was discontinued,[11, 13] we also found that the initial albuminuria response significantly correlated with the response after drug discontinuation, although the correlation was weaker compared to the studies with shorter follow-up.

Correlation in response after re-exposure

Thirdly, to further explore if the between individual variability in response is related or unrelated to drug treatment, we analyzed a cross-over study in which 17 patients were exposed twice to the same intervention and calculated how much of the between patient variability at second exposure could be explained by the first exposure.[19] We observed that 53% and 63% of the between-individual variability in albuminuria following the second exposure to lisinopril 10 mg/d or losartan 50 mg/d was explained by the first exposure to these agents, respectively (Figure 3). The positive predictive value and negative predictive value of the first albuminuria response to predict a 40% reduction in albuminuria to losartan or lisinopril during the second exposure were 72.7% and 71.4%, and the area under the ROC curve was 0.84. The positive predictive value, negative predictive value and area under the ROC curve to predict a 50% albuminuria reduction during re-exposure were 83.3%, 75.0%, and 0.78, respectively.

Finally, the correlation between the initial and post-treatment albuminuria change, as well as the correlation in response after re-exposure were consistent regardless which drugs were used.

Discussion

Individual patients show a large variability in their albuminuria response after the initiation of treatment. Whether this variability represents a true response variability or a random variability unrelated to treatment was not previously studied. In this analysis of multiple trials determining the albuminuria-lowering effects of different classes of drugs in patients with and without diabetes we can detect a true marked response variation in albuminuria lowering to varying drugs between patients, despite within-individual fluctuations in albuminuria over time of around 25%. This variation is true as supported by several results: first, 50 to 60% of the variation after re-exposure to the same drug in the same patient can be explained by the variation during the first exposure. Second, the initial degree of albuminuria response appears to correlate with the change in albuminuria after drug discontinuation, even after several years albuminuria returns to baseline values again.

Two previous rotation studies have drawn similar conclusions using blood pressure instead of albuminuria data: blood pressure measurements fluctuate within an individual, blood pressure responses to anti-hypertensive drugs vary consistently between individuals, as evidenced by the finding that re-exposure to anti-hypertensive medication results in a similar blood pressure response at re-exposure.[20-22] An additional finding in these rotation studies was that patients who responded well to one class of drug responded poorly to another class of drug. This led to the Cambridge AB/CD rule for optimization of antihypertensive drugs because the individual responses to ACE-inhibitors and B-blockers correlated with each other and so did the responses to Calcium Antagonist and Diuretics. We did see a clear difference in the average albuminuria lowering response to the different drugs tested in our different studies. However, since we did not expose each patient to multiple of these drugs, we cannot conclude whether there is an optimal drug for each patient.

A recent meta-analysis concluded that the evidence of between patient variability in blood pressure responses to ACE-inhibitors is weak.[10] This meta-analysis included multi-national trials not designed to analyze between-individual drug response variability in contrast to prior prospectively designed rotation blood pressure trials.[20] Various factors such as therapy adherence, changes in background medication, and disease progression influence individual drug response. These factors may have biased the conclusion that most of the response variation to antihypertensive medication is due to within-individual day-to-day fluctuations in blood pressure.

What could explain the individual drug response variability? It is possible that the individual response in albuminuria is explained by individual blood pressure responses. However, this possibility is unlikely since previous studies have shown that the blood pressure and albuminuria response to RAS intervention within an individual are dissociated.[23, 24] In other words, some patients show a reduction in both blood pressure and albuminuria, but in other patients only blood pressure or albuminuria is reduced. Indeed, adjustment for individual changes in blood pressure in the current study did not alter our findings. Various other factors may be involved in individual differences in responsiveness to RAS-intervention, such as genetic factors,[25] extracellular volume status,[26] or the type and severity of renal disease. Using novel 'omic' and systems medicine network approaches provide new opportunities to understand the underlying determinants of drug response variability.[27] Additionally, imaging techniques to better characterize and visualize drug disposition in vivo may also provide new avenues to study determinants of drug response variability.

Prior studies have already shown that albuminuria shows substantial day-to-day variability.[28] We confirmed these findings and also showed in a large cohort that the individual day-to-day variability is consistent over time. In addition, we showed that the variability depends on the baseline albuminuria level such that the largest intra-individual albuminuria variability over time is observed in patients with microalbuminuria while within-individual variations in albuminuria are smaller in the low or high albuminuria range.

The practical implication of our study is that clinicians should measure and monitor the initial response of each individual to albuminuria lowering intervention. The alternative of taking the average drug response observed in clinical trials is not recommended since a substantial part of the total between-individual variability in response could be attributed to drug response variability. If the reduction in albuminuria following intervention is insufficient, up-titrating the dose of the drug, moderation of dietary sodium intake, or adding diuretic treatment to the therapeutic regimen may further lower albuminuria and potentially improve prognosis.[29-31]

This study has limitations. First of all we used data from several trials that were not designed to address our research question. Secondly, we were unable to include all clinical studies with an albuminuria treatment response and recovery period in our analysis due to inaccessibility of individual patient data. In addition, only one trial could be included that exposed the same individual twice to the same drug to assess whether individual drug responses on albuminuria are consistent. An ongoing trial (IMPROVE; Dutch trial registry NTR4439) is designed to prospectively study between-individual variability in drug response in albuminuria by multiple exposures. Another limitation is that plasma drug levels were not available in four studies to verify that patients adhered to study medication.

In conclusion, the variable changes in albuminuria following drug initiation can be separated from the day-to-day within-individual variability in albuminuria, indicating that individual drug responses can be adequately quantified in a clinical trial setting. Monitoring for albuminuria responses is thus feasible and recommended in practice. The factors that determine drug response should be investigated in future studies, in order to develop measures to improve drug response in non-responsive patients. This is of particular relevance since the initial effect of a drug on albuminuria is related to its long term renal protective effect.

What is already known about this subject?

Individual patients show a wide range of responses to albuminuria lowering drugs. Since albuminuria varies considerably from day-to-day we questioned whether this variability in response reflects a random variability or a true drug response.

What this study adds

This study showed that the variable changes in albuminuria after start of Albuminuria lowering drugs can be separated from day-to-day fluctuations in albuminuria indicating that individual drug responses can be adequately quantified in clinical trial settings. These data support further research into the factors that determine drug response in order to develop measures to overcome therapy resistance to albuminuria lowering drugs. This is relevant since the initial effect of a drug on albuminuria is related to its long term renal protective effect.

Acknowledgements

The work described in this paper received funding from the Novo Nordisk Foundation, Grant Number NNF14SA0003. HJL Heerspink is supported by a VIDI grant from the Netherlands Organisation for Scientific Research (917.15.306).

Conflicts of interest

SP reports no conflicts of interest. DdZ is consultant for and received honoraria (to employer) from AbbVie, Astellas, Eli-Lilly, Chemocentryx, Fresenius, and Janssen. FP and PR are employed at Steno Diabetes Center, Gentofte, Denmark. Steno Diabetes Center is an independent academic institution owned by Novo Nordisk and The Novo Nordisk Foundation. FP reports having received research grants from Astra Zeneca, lecture fees from Astra Zeneca, MSD, Janssen, Lily, Boehringer Ingelheim, Novo Nordisk, Novartis and being consultant/advisory board member for Astra Zeneca and MSD. PR received lecture fees from Novartis and Boehringer Ingelheim, and research grant from Novartis, Astra Zeneca, Novo Nordisk and has served as a consultant for Merck, Astra Zeneca, Boehringer Ingelheim, MSD, AbbVie, Novo Nordisk and having equity interest in Novo Nordisk. RTG is consultant and/or received honoraria (to employer) from Abbvie, Bayer, Genzyme, Ipsen, and Otsuka. HJLH is consultant for and received honoraria (to employer) from AbbVie, Astellas, Astra Zeneca, Boehringer Ingelheim, Janssen, Merck, and ZS-Pharma .

References

1. Schievink B, de Zeeuw D, Parving HH, Rossing P, Lambers Heerspink HJ: The renal protective effect of Angiotensin Receptor Blockers depends on intra-individual response variation in multiple risk markers. *Br J Clin Pharmacol* 2015
2. Heerspink HJ, Kropelin TF, Hoekman J, de Zeeuw D: Drug-Induced Reduction in Albuminuria Is Associated with Subsequent Renoprotection: A Meta-Analysis. *J Am Soc Nephrol* 2015 **26**:2055-64
3. Lambers Heerspink HJ, Gansevoort RT: Albuminuria Is an Appropriate Therapeutic Target in Patients with CKD: The Pro View. *Clin J Am Soc Nephrol* 2015 **10**:1079-88
4. Poortmans J, Dorchy H, Toussaint D: Urinary excretion of total proteins, albumin, and beta 2-microglobulin during rest and exercise in diabetic adolescents with and without retinopathy. *Diabetes Care* 1982 **5**:617-23
5. Clerico A, Giammattei C, Cecchini L, Lucchetti A, Cruschelli L, Penno G, Gregori G, Giampietro O: Exercise-induced proteinuria in well-trained athletes. *Clin Chem* 1990 **36**:562-4
6. Brinkman JW, de Zeeuw D, Duker JJ, Gansevoort RT, Kema IP, Hillege HL, de Jong PE, Bakker SJ: Falsely low urinary albumin concentrations after prolonged frozen storage of urine samples. *Clin Chem* 2005 **51**:2181-3
7. Wagner MG, Smith FG, Jr., Tinglof BO, Jr., Cornberg E: Epidemiology of proteinuria. A study of 4,807 schoolchildren. *J Pediatr* 1968 **73**:825-32
8. Rytand DA, Spreiter S: Prognosis in postural (orthostatic) proteinuria: forty to fifty-year follow-up of six patients after diagnosis by Thomas Addis. *N Engl J Med* 1981 **305**:618-21
9. Hemmingsen L, Skaarup P: Urinary excretion of ten plasma proteins in patients with febrile diseases. *Acta Med Scand* 1977 **201**:359-64
10. Bell KJ, Hayen A, Macaskill P, Craig JC, Neal BC, Fox KM, Remme WJ, Asselbergs FW, van Gilst WH, Macmahon S, Remuzzi G, Ruggenti P, Teo KK, Irwig L: Monitoring initial response to Angiotensin-converting enzyme inhibitor-based regimens: an individual patient data meta-analysis from randomized, placebo-controlled trials. *Hypertension* 2010 **56**:533-9
11. Asselbergs FW, Diercks GF, Hillege HL, van Boven AJ, Janssen WM, Voors AA, de Zeeuw D, de Jong PE, van Veldhuisen DJ, van Gilst WH: Effects of fosinopril and

- pravastatin on cardiovascular events in subjects with microalbuminuria. *Circulation* 2004 **110**:2809-16
12. Ruggenti P, Fassi A, Ilieva AP, Bruno S, Iliev IP, Brusegan V, Rubis N, Gherardi G, Arnoldi F, Ganeva M, Ene-Iordache B, Gaspari F, Perna A, Bossi A, Trevisan R, Dodesini AR, Remuzzi G: Preventing microalbuminuria in type 2 diabetes. *N Engl J Med* 2004 **351**:1941-51
 13. Parving HH, Lehnert H, Brochner-Mortensen J, Gomis R, Andersen S, Arner P: The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *N Engl J Med* 2001 **345**:870-8
 14. Brenner BM, Cooper ME, de Zeeuw D, Keane WF, Mitch WE, Parving HH, Remuzzi G, Snapinn SM, Zhang Z, Shahinfar S: Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001 **345**:861-9
 15. Gansevoort RT, de Zeeuw D, de Jong PE: Is the antiproteinuric effect of ACE inhibition mediated by interference in the renin-angiotensin system? *Kidney Int* 1994 **45**:861-7
 16. Persson F, Rossing P, Reinhard H, Juhl T, Stehouwer CD, Schalkwijk C, Danser AH, Boomsma F, Frandsen E, Parving HH: Optimal antiproteinuric dose of aliskiren in type 2 diabetes mellitus: a randomised crossover trial. *Diabetologia* 2010 **53**:1576-80
 17. de Zeeuw D, Coll B, Andress D, Brennan JJ, Tang H, Houser M, Correa-Rotter R, Kohan D, Lambers Heerspink HJ, Makino H, Perkovic V, Pritchett Y, Remuzzi G, Tobe SW, Toto R, Viberti G, Parving HH: The Endothelin Antagonist Atrasentan Lowers Residual Albuminuria in Patients with Type 2 Diabetic Nephropathy. *J Am Soc Nephrol* 2014;**25**:1083-93
 18. de Zeeuw D, Agarwal R, Amdahl M, Audhya P, Coyne D, Garimella T, Parving HH, Pritchett Y, Remuzzi G, Ritz E, Andress D: Selective vitamin D receptor activation with paricalcitol for reduction of albuminuria in patients with type 2 diabetes (VITAL study): a randomised controlled trial. *Lancet* 2010 **376**:1543-51
 19. Laverman GD, Henning RH, de Jong PE, Navis G, de Zeeuw D: Optimal antiproteinuric dose of losartan in nondiabetic patients with nephrotic range proteinuria. *Am J Kidney Dis* 2001 **38**:1381-4
 20. Dickerson JE, Hingorani AD, Ashby MJ, Palmer CR, Brown MJ: Optimisation of antihypertensive treatment by crossover rotation of four major classes. *Lancet* 1999 **353**:2008-13

21. Attwood S, Bird R, Burch K, Casadei B, Coats A, Conway J, Dawes M, Ebbs D, Farmer A, Robinson J: Within-patient correlation between the antihypertensive effects of atenolol, lisinopril and nifedipine. *J Hypertens* 1994 **12**:1053-60
22. Williams B, MacDonald TM, Morant S, Webb DJ, Sever P, McInnes G, Ford I, Cruickshank JK, Caulfield MJ, Salsbury J, Mackenzie I, Padmanabhan S, Brown MJ: Spironolactone versus placebo, bisoprolol, and doxazosin to determine the optimal treatment for drug-resistant hypertension (PATHWAY-2): a randomised, double-blind, crossover trial. *Lancet* 2015 **386**:2059-68
23. Hellemons ME, Persson F, Bakker SJ, Rossing P, Parving HH, De Zeeuw D, Lambers Heerspink HJ: initial angiotensin receptor blockade-induced decrease in albuminuria is associated with long-term renal outcome in type 2 diabetic patients with microalbuminuria: a post hoc analysis of the IRMA-2 trial. *Diabetes Care* 2012 **34**:2078-83
24. Eijkelkamp WB, Zhang Z, Remuzzi G, Parving HH, Cooper ME, Keane WF, Shahinfar S, Gleim GW, Weir MR, Brenner BM, de Zeeuw D: Albuminuria is a target for renoprotective therapy independent from blood pressure in patients with type 2 diabetic nephropathy: post hoc analysis from the Reduction of Endpoints in NIDDM with the Angiotensin II Antagonist Losartan (RENAAL) trial. *J Am Soc Nephrol* 2007 **18**:1540-6
25. Parving HH, de Zeeuw D, Cooper ME, Remuzzi G, Liu N, Luncford J, Shahinfar S, Wong PH, Lyle PA, Rossing P, Brenner BM: ACE gene polymorphism and losartan treatment in type 2 diabetic patients with nephropathy. *J Am Soc Nephrol* 2008 **19**:771-9
26. Slagman MC, Waanders F, Hemmelder MH, Woittiez AJ, Janssen WM, Lambers Heerspink HJ, Navis G, Laverman GD: Moderate dietary sodium restriction added to angiotensin converting enzyme inhibition compared with dual blockade in lowering proteinuria and blood pressure: randomised controlled trial. *Bmj* 2011 **343**:d4366
27. Lambers Heerspink HJ, Oberbauer R, Perco P, Heinzl A, Heinze G, Mayer G, Mayer B: Drugs meeting the molecular basis of diabetic kidney disease: bridging from molecular mechanism to personalized medicine. *Nephrol Dial Transplant* 2015 **30** Suppl 4:iv105-12
28. Witte EC, Lambers Heerspink HJ, de Zeeuw D, Bakker SJ, de Jong PE, Gansevoort R: First morning voids are more reliable than spot urine samples to assess microalbuminuria. *J Am Soc Nephrol* 2009 **20**:436-43

29. Burgess E, Muirhead N, Rene de Cotret P, Chiu A, Pichette V, Tobe S: Supramaximal dose of candesartan in proteinuric renal disease. *J Am Soc Nephrol* 2009 **20**:893-900
30. Heerspink HJ, Holtkamp FA, Parving HH, Navis GJ, Lewis JB, Ritz E, de Graeff PA, de Zeeuw D: Moderation of dietary sodium potentiates the renal and cardiovascular protective effects of angiotensin receptor blockers. *Kidney Int* 2012 **82**:330-7
31. Vogt L, Waanders F, Boomsma F, de Zeeuw D, Navis G: Effects of dietary sodium and hydrochlorothiazide on the antiproteinuric efficacy of losartan. *J Am Soc Nephrol* 2008 **19**:999-1007

Figure 1




Figure 1: Within individual coefficient of variation higher in patients with micro- or macroalbuminuria compared to normoalbuminuria. * $P < 0.01$

Accepted

Figure 2




Figure 2: Correlation between the initial albuminuria response and response after drug discontinuation across trials.

Footnote: *Atrasentan 0.75mg; **Atrasentan 1.25 mg; †Aliskiren 150mg; ‡Aliskiren 300 mg; §Aliskiren 600 mg

Accepted

Figure 3




Figure 3: Albuminuria responses when the same individual is exposed twice to the same drug at the same dose

Accepted

Table 1: Within-individual coefficient of variation of albuminuria at different time-intervals in the placebo arms of trials in non-diabetic and diabetic patients with high normo-, micro- or macroalbuminuria.

Trial	N	Baseline albuminuria	Time- interval	CV (25th – 75th Percentile)
PREVEND-IT	127	23	0-12-24	26.9 [16.2 – 45.5]
			24-36-48	25.0 [16.3 – 40.0]
BENEDICT	120	5.8	0-6-12	22.2 [13.6 – 39.6]
			12-18-24	20.0 [14.1 – 32.4]
			24-30-36	24.0 [17.6 – 43.3]
IRMA-2	125	53	0-6-12	35.7 [22.8 – 56.5]
			12-18-24	37.7 [26.5 – 57.5]
RENAAL	205	1013	0-6-12	34.9 [20.6 – 54.0]
			12-18-24	33.4 [20.5 – 50.4]
			24-30-36	35.5 [22.7 – 56.6]

Table 2: Study characteristics

	Gansevoort N=11	Persson (N=19)	RADAR (N=174)	VITAL (N=140)	IRMA-2 (N=73)	PREVEND-IT (N=655)
Age, years	42.2 (12)	64.3 (7)	64.7 (9)	64.4 (10)	56.1 (9)	51.2 (11)
Female, N (%)	2 (18.2)	1 (5)	42 (24)	46 (33)	20 (27.4)	216 (33)
Albuminuria (e)GFR (ml/min/1.73m ²)	4.6*	553 [180 – 746]	827 [465 – 1516]	567 [261-1114]	50 [30 – 80]	23 [16 – 43]
Systolic BP (mmHg)	70.7 (17)	80.4 (23)	49.4 (14)	42.0 (18)	75.3 (13)	75.7 (12)
Diabetes, N (%)	153 (11)	136.6 (10)	136.5 (14)	140.9 (16)	151 (12)	130.2 (17)
Intervention	0 (0)	19 (100)	174 (100)	140 (100)	73 (100)	0 (0)
Follow-up duration	ACEi / ARB (Enalapril 20mg / Losartan 100 mg)	DRI (Aliskiren 150, 300, 600mg)	ERA (Atrasentan 0.75, 1.25mg)	VDRA (Paricalcitol 2µg)	ARB (Irbesartan 150/300 mg)	ACEi (Fosinopril 20mg)
	4 weeks	4 weeks	12 weeks	24 weeks	2 years	4 years

* Proteinuria in 24-hr urine collection; ACEi, Angiotensin Converting Enzyme Inhibitor; ARB, Angiotensin Receptor Blocker, ERA, Endothelin Receptor Antagonist; VDRA, Vitamin D Receptor Antagonist; ARB, Angiotensin Receptor Blocker; ACEi, Angiotensin Converting Enzyme Inhibitor

Table 3: Albuminuria response after start treatment (on-drug response), albuminuria response after drug discontinuation (off-drug response), and correlation between on-drug and off-drug response within each trial

Correlation on and off drug response	N	Mean drug response after start and discontinuation						Correlations in on-drug and off-drug response within each trial			
		On-drug albuminuria response (%)	P25*	P75	Off-drug albuminuria response (%)	P25	P75	Regression Coefficient	P-value	R2	r
<i>Macroalbuminuria: Gansevoort</i>											
Losartan 100 mg	11	-49.2	-61.5	-20.9	39.3	31.1	47.4	-0.47	0.015	0.465	0.682
Enalapril 20 mg	11	-56.8	-69.2	-29.0	47.5	17.5	71.4	-0.79	0.007	0.569	0.754
<i>Macroalbuminuria: Persson</i>											
Placebo	19	-8.8	-44.6	32.9	-0.4	-40.0	19.3	-0.35	0.111	0.143	0.377
Aliskiren 150 mg	19	-42.7	-61.9	-28.4	36.9	12.0	56.5	-0.10	0.675	0.011	0.103
Aliskiren 300 mg	19	-54.6	-73.1	-7.1	50.0	14.0	68.5	-0.49	0.005	0.375	0.612
Aliskiren 600 mg	19	-59.4	-79.5	-21.4	55.4	35.6	77.8	-0.55	0.001	0.472	0.687
<i>Macroalbuminuria: RADAR</i>											
Placebo	46	2.8	-18.2	29.6	-4.5	-26.6	16.1	-0.08	0.514	0.009	0.095
Atrasentan 0.75 mg	64	-35.9	-51.2	-4.0	40.4	14.2	58.4	-0.44	<0.001	0.268	0.518
Atrasentan 1.25 mg	64	-40.7	-56.4	-15.9	36.6	10.9	49.1	-0.38	<0.001	0.142	0.377
<i>Macroalbuminuria: VITAL</i>											
Placebo	71	-7.1	-37.0	38.2	-9.1	-44.3	21.5	-0.03	0.77	0.001	0.035
Paricalcitol 2 µg	64	-18.8	-49.4	17.2	13.3	-19.7	40.5	-0.04	0.69	0.002	0.049
<i>Normoalbuminuria: PREVEND-IT</i>											
Placebo	325	0.1	-21.0	29.0	1.2	-34.6	24.6	-0.04	0.574	0.001	0.032
Fosinopril 20 mg	330	-24.3	-45.7	2.0	23.5	-15.7	46.0	-0.13	0.025	0.015	0.122
<i>Microalbuminuria: IRMA-2</i>											
Placebo	29	20.2	0.0	60.8	11.8	-16.7	39.6	0.25	0.217	0.056	0.236
Irbesartan 150/300 mg	47	-37.3	-56.0	0.0	30.5	0.0	59.5	-0.47	0.046	0.085	0.292

*P25/P75 denotes the first and third quartile of the on-drug and off-drug response